

Title	Phylum Arthropoda von Siebold, 1848. <i>In</i>: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness
Authors	ZHI-QIANG ZHANG
Publication date	2011/12/23
Journal name	Zootaxa
Volume	3148
Pages	99–103
Description	A new evolutionary classification of the phylum Arthropoda is presented. The survey provides the most updated count of 1,242,040 species described from the world.

Phylum **Arthropoda** von Siebold, 1848¹ (1,242,040 species, of which †27,745)²
 Subphylum 1 †**Trilobitomorpha** Størmer, 1944 (19,606 species)³
 Subphylum 2 **Chelicerata** Heymons, 1901 (113,894 species, of which 1,957)⁴
 Class 1 **Pycnogonida** Latreille, 1810 (1330 species, of which †8 species)⁵
 Class 2 †**Aglaspidida** Walcott, 1911 (†11 species)⁶
 Class 3.1 †**Xiphosura** Latreille, 1802 (†98 species)⁷
 Class 3.2.1 †**Eurypterida** Burmeister, 1843 (†246 species)⁸
 Class 3.2.2 †**Chasmataspida** Caster & Brooks, 1956 (†8 species)⁹
 Class 3.3 **Arachnida** Cuvier, 1812 (112,201, of which †1,586)¹⁰
 Order 1.1 **Opiliones** Sundevall, 1833 (6,519 species, of which †35 species)¹¹
 Order 1.2 **Scorpiones** C.L. Koch, 1851 (2,068 species, of which †121 species)¹²
 Order 2.1 **Solifugae** Sundevall, 1833 (1,116 species, of which †3 species)¹³
 Order 2.2 **Pseudoscorpiones** de Geer, 1778 (3,494 species, of which †40 species)¹⁴
 Order 3 **Palpigradi** Thorell, 1888 (83 species, of which †1 species)¹⁵
 Order 4 †**Phalangiotarbida** Haase, 1890 (†31 species)¹⁶
 Order 5 **Ricinulei** Thorell, 1876 (73 species, of which †15 species)¹⁷
 Order 6.1 **Opilioacarida** Zakhvatkin, 1952 (37 species, of which †2 species)¹⁸
 Order 6.2 **Holothyrida** Thon, 1905 (27 species)
 Order 6.3 **Ixodida** Leach, 1815 (896 species, of which †5 species)¹⁹
 Order 6.4 **Mesostigmata** G. Canestrini, 1891 (11,424 species)
 Order 7.1 **Trombidiformes** Reuter, 1909 (25,821 species, of which †24 species)²⁰
 Order 7.2 **Sarcoptiformes** Reuter, 1909 (16,412 species, of which †113 species)²¹
 Order 8.1 †**Trigonotarbita** Petrunkevitch, 1949 (†65 species)²²
 Order 8.2.1.1 †**Uraraneida** Selden & Shear in Selden *et al.*, 2008 (†2 species)²³
 Order 8.2.1.2 **Araneae** Clerck, 1757 (43,579 species, of which †1,106 species)²⁴
 Order 8.2.2.1 †**Haptopoda** Pocock, 1911 (†1 species)²⁵

1. **BY** Zhi-Qiang Zhang (for full author addresses, see **Author name and address** after **References**). The title of this contribution should be cited as “Phylum Arthropoda von Siebold, 1848 *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness”. This classification at the subphylum level generally reflects the high level relationship summarized by Giribet & Edgecombe (2012: Fig. 5), with modification and updates as indicated (see notes on Crustacea).
 Fossil Marellomorpha not included in the phylogenetic relationships focused on Recent taxa in Giribet & Edgecombe (2012). Grimaldi & Engel (2005) listed three genera.
2. Only Recent species were counted for many insect orders and all of Myriapoda; so the total number of species should be considered incomplete, as diversity of fossil taxa is underestimated.
3. See Adrain (2011, this volume).
4. There are broad consensus of high level relationship within Chelicerata (Dunlop 2010; Giribet & Edgecombe 2011); this classification at the class level generally reflects the high level relationship summarized by Giribet & Edgecombe (2011: Fig. 5), with modification and updates as indicated.
 Grimaldi & Engel (2005) listed three poorly known arthropod taxa (†Sidneyida, †Emeraldellida Størmer, 1944 and †Sanctacarida); their placement is uncertain in the current system (e.g. Dunlop 2010; Giribet & Edgecombe 2012).
5. See Bamber (2011, this volume).
6. Data from J. Ortega-Hernandez (personal communication, 15 Dec. 2011).
7. Fossil horseshoe crabs; based on Dunlop *et al.* (2011).
8. Fossil sea scorpions; based on Dunlop *et al.* (2011).
9. Based on Dunlop *et al.* (2011).
10. Arachnid classification reflects mostly the phylogenetic hypothesis by Shultz (2007), but his Acaromorpha is now disputed (see Dunlop 2010).
11. See Kury (2011, this volume).
12. See Prendini (2011a, this volume).
13. See Prendini (2011b, this volume).
14. See Harvey (2011, this volume).
15. See Prendini (2011c, this volume).
16. Based on Dunlop *et al.* (2011).
17. See Prendini (2011d, this volume).
18. See Beaulieu *et al.* (2011, this volume), with a list of families and diversity estimates for each family within the mite superorder Parasitiformes, including Opilioacarida, Holothyrida, Ixodida and Mesostigmata.
19. Based on Guglielmone *et al.* (2010).
20. See Zhang *et al.* (2011, this volume). The second mite superorder, Acariformes, consists of Trombidiformes and Sarcoptiformes.
21. This order includes two suborders: Oribatida and Endeostigmata. According to Schatz *et al.* (2011, this volume), Recent Oribatida (including Astigmata) currently include 249 families, 2399 genera and 16197 species; exclusively fossil taxa include 2 families, 20 genera and 108 species; Walter *et al.* (2011, this volume) estimated 10 families, 27 genera, 108 species for Endeostigmata, of which 3 genera and 5 species are based on fossil.
22. Based on Dunlop *et al.* (2011).
23. Selden *et al.* (2008).
24. See Dunlop & Penney (2011, this volume).

- Order 8.2.2.2.1 **Amblypygi** Thorell, 1883 (170 species, of which †9 species)²⁶
 Order 8.2.2.2.2.1 **Thelyphonida** Latreille, 1804 (119 species, of which †9 species)²⁷
 Order 8.2.2.2.2.2 **Schizomida** Petrunkevitch, 1945 (264 species, of which †4 species)²⁸
 Subphylum 2.1 **Myriapoda** Latreille, 1802 (11,885 species)²⁹
 Class 1 **Chilopoda** Latreille, 1817 (3,100 species)³⁰
 Class 2.1 **Symphyla** Ryder, 1880 (197 species)³¹
 Class 2.2.1 **Pauropoda** Lubbock, 1868 (835 species)
 Class 2.2.2 **Diplopoda** de Blainville in Gervais, 1844 (7,753 species)
 Subphylum 2.2.1 **Crustacea** Brünnich, 1772 (66,914 species)³²
 Subphylum 2.2.2 **Hexapoda** Latreille, 1825 (1,029,741 species, of which †6,182 species)³³
 Class 1.1 **Collembola** Lubbock, 1870 (8,130 species)³⁴
 Class 1.2.1 **Protura** Silvestri, 1907 (804 species)³⁵
 Class 1.2.2 **Diplura** Börner, 1904 (800 species)³⁶
 Class 2 **Insecta** Linnaeus, 1758 (1,020,007 species, of which †6,182 species)³⁷
 Order 1 **Archaeognatha** (513 species)³⁸
 Order 2.1 **Zygentoma** Börner, 1904 (561 species, of which †1 species)³⁹
 Order 2.2.1.1 **Ephemeroptera** Hyatt & Arms, 1890 (3,240 species)⁴⁰
 Order 2.2.1.2.1.1 †**Geroptera** Brodsky, 1994 (†2 species)⁴¹
 Order 2.2.1.2.1.2.1 †**Protodonata** (†42 species)⁴²
 Order 2.2.1.2.1.2.2 **Odonata** Fabricius, 1792 (5,899 species)⁴³
 Order 2.2.1.2.2.1.1 †**Palaeodictyoptera** Goldenberg, 1877 (†115 species)⁴⁴
 Order 2.2.1.2.2.2.1 **Mischopterida** Handlirsch, 1906 (†100 species)⁴⁵
 Order 2.2.1.2.2.2.2 †**Diaphanopteroidea** Handlirsch, 1906 (†50 species)⁴⁶
 Order 2.2.2.1 †**Paoliida** Handlirsch, 1906 (†12 species)⁴⁷
 Order 2.2.2.2.1 †**Caloneuroidea** Martynov, 1938 (†14 species)⁴⁸
 Order 2.2.2.2.2 †**Titanoptera** Sharov, 1968 (†15 species)⁴⁹

25. Based on Dunlop *et al.* (2011).
 26. See Prendini (2011e, this volume).
 27. See Prendini (2011f, this volume).
 28. See Prendini (2011g, this volume).
 29. Recent species only; many fossil species were described but were not included in this edition due to lack of time, so the total number of species should be considered incomplete.
 30. See Shear (2011, this volume).
 31. See Minelli (2011, this volume), which also covers Pauropoda and Diplopoda.
 32. See Ahyong *et al.* (2011, this volume), which provided a classification of all families and an estimate of 1,003 families, 9,522 genera and 66,914 species for Recent Crustacea. Crustacea is paraphyletic according to Giribet & Edgecombe (2011). Here traditional consensus classification presented in Ahyong *et al.* (2011) is followed.
 33. Classification reflects consensus phylogenetic relationship in Trautwein *et al.* (2012); inclusion of fossil orders follows Grimaldi & Engel (2005).
 34. See Janssens & Christiansen (2011, this volume).
 35. Based on Szeptycki (2007), and updated using Zoological Records from 2007.
 36. This number is cited by Chapman (2009) and also Tree of Life Web Project. 1995. Diplura. Version 01 January 1995 (temporary). <http://tolweb.org/Diplura/8204/1995.01.01> in The Tree of Life Web Project, <http://tolweb.org/>.
 37. Diversity estimates for fossil insects are not available for many orders and thus total diversity is underestimated.
 38. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 39. Alternative name Thysanura; estimates based Foottit & Adler (2009) and updated using Zoological Records from 2008. Fossil taxa are represented in at least 1 family (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data may be incomplete.
 40. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008; Foottit & Adler listed 3,046 species.
 41. Fossil taxa are represented in 2 genera and 1 family (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data may be incomplete; Grimaldi & Engel (2005) mentioned “a few species” for this order.
 42. Fossil taxa are represented in 27 genera and 5 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete. Zoological Record lists a total of 48 species/subspecies names.
 43. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008; Foottit & Adler listed 5,680 species; Trueman (2007) mentioned “around 6,000 species” for Odonata.
 44. Fossil taxa are represented in at least 82 genera and 25 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete.
 45. Concept and estimates follow Rasnitsyn & Quicke (2002); this order includes two orders listed in Grimaldi & Engel (2005): †Dipliptera (=Archodonata Martynov, 1932) and †Megasecoptera Brongniart, 1885.
 46. Based on Rasnitsyn & Quicke (2002).
 47. Based on Prokop & Nel (2007).
 48. Fossil taxa are represented in at least 11 genera and 7 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete; a total of 25 genera 9 families in David Eades. Polyneoptera Species File Online. Version 1.0/4.0 [retrieval date]. <http://Polyneoptera.SpeciesFile.org>, but Eades did not provide species for some genera.

- Order 2.2.2.2.3 **Orthoptera** Olivier, 1789 (24,276 species, of which †421 species)⁵⁰
 Order 2.2.2.3.1 **Phasmida** Leach, 1815 (3,029 species, of which †15 species)⁵¹
 Order 2.2.2.3.2 **Embioptera** Lameere, 1900 (464 species⁵², of which †1 species)
 Order 2.2.2.4.1 **Grylloblattodea** Brues & Melander, 1932 (34 species)⁵³
 Order 2.2.2.4.2 **Mantophasmatodea** Zompro, Klass, Kristensen & Adis, 2002 (21 species, of which †6 species)⁵⁴
 Order 2.2.2.5.1 **Plecoptera** Latreille 1802 (3,788 species⁵⁵, of which †45 species)⁵⁶
 Order 2.2.2.5.2 **Dermaptera** deGeer, 1773 (1,978 species)⁵⁷
 Order 2.2.2.6.1 **Zoraptera** Silvestri, 1913 (37 species)⁵⁸
 Order 2.2.2.6.2.1 **Mantodea** Burmeister 1838 (2,400 species)⁵⁹
 Order 2.2.2.6.2.2 **Blattodea** Brunner, 1882 (7,314 species)⁶⁰
 Order 2.2.3.1 †**Miomoptera** Martynov 1927 (†47 species)⁶¹
 Order 2.2.3.2.1 **Psocoptera** Shipley, 1904 (5,720 species)⁶²
 Order 2.2.3.2.2 **Phthiraptera** Haeckel, 1896 (5,102 species)⁶³
 Order 2.2.3.3.1 **Thysanoptera** Haliday, 1836 (6,019 species, of which †155 species)⁶⁴
 Order 2.2.3.3.2 **Hemiptera** Linnaeus 1758 (103,590)⁶⁵
 Order 2.2.4 †**Glosselytrodea** Martynov 1938 (†30 species)⁶⁶
 Order 2.2.5.1 **Hymenoptera** Linnaeus 1758⁶⁷ (116,861 species)⁶⁸
 Order 2.2.5.2.1.1.1 **Strepsiptera** Kirby 1813 (609 species)⁶⁹
 Order 2.2.5.2.1.1.2 **Coleoptera** Linnaeus, 1758 (387,100 species, of which †600 species)⁷⁰
 Order 2.2.5.2.1.2.1 **Neuroptera** Linnaeus, 1758 (5,868 species)⁷¹
 Order 2.2.5.2.1.2.2.1 **Megaloptera** Latreille, 1802 (354 species)⁷²
 Order 2.2.5.2.1.2.2.2 **Raphidioptera** Navás, 1916 (254 species)⁷³
 Order 2.2.5.2.2.1.1 **Trichoptera** Kirby, 1813 (14,999 species, of which †608 species)⁷⁴
 Order 2.2.5.2.2.1.2 **Lepidoptera** Linnaeus, 1758 (157,424 species, of which †86 species)⁷⁵
 Order 2.2.5.2.2.2.1 **Diptera** Linnaeus, 1758 (159,294 species, of which †3,817 species)⁷⁶
 Order 2.2.5.2.2.2.2.1 **Siphonaptera** Latreille, 1825 (2,075 species)⁷⁷
 Order 2.2.5.2.2.2.2.2 **Mecoptera** Packard, 1886 (757 species)⁷⁸

49. Based on Shcherbakov (2011).
 50. See Ingrisch (2011, this volume).
 51. See Brock and Marshall (2011, this volume) for Recent taxa only. Fossil taxa are represented in at least 14 genera and 9 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete.
 52. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 53. Grimaldi & Engel (2005) mentioned 26 species placed in 5 genera in 1 family.
 54. Updated with Arillo & Engel (2006), Eberhard *et al.* (2011). Note that this order and Grylloblattodea were placed in the order Notoptera by Arillo & Engel (2006).
 55. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008; Foottit & Adler listed 3497 species.
 56. Fossil taxa are represented in at least 22 genera and 10 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete.
 57. Based on Foottit & Adler (2009) and updated using Zoological Records.
 58. Based on Foottit & Adler (2009) and updated using Zoological Records.
 59. Based on Otte, D., Spearman, L. & Martin, B.D.S. Mantodea Species File Online. Version 1.0/4.0. [retrieval date 15 Dec. 2011]. <<http://Mantodea.SpeciesFile.org>>.
 60. See Beccaloni & Eggleton (2011, this volume), without fossils; including 2,692 species of termites (no longer an order).
 61. Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search; this should be viewed as incomplete.
 62. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 63. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 64. See Mound *et al.* (2011, this volume).
 65. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 66. Based on Hong (2007) with updates.
 67. Holometabola for 11 orders from Hymenoptera to Mecoptera.
 68. Based on Hymenoptera Online <http://hol.osu.edu/> (accessed 15 Dec. 2011), which placed these species in 7,745 genera and 118 families. Fossil taxa are represented in at least 789 genera and 89 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete. Foottit & Adler (2009), citing Huber in that volume, mentioned 144,695 species.
 69. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 70. See Slipinski *et al.* (2011, this volume).
 71. Based on Foottit & Adler (2009) and updated using Zoological Records.
 72. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008; Foottit & Adler listed 328 species.
 73. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 74. See Holzenthall *et al.* (2011, this volume).
 75. See Nieuwerkerken *et al.* (2011, this volume).
 76. See Pape *et al.* (2011, this volume), but 5,969 dubious species not included in total count.
 77. Based on Foottit & Adler (2009) and updated using Zoological Records.
 78. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.

Acknowledgements

I thank Dr Rich Leschen (Landcare Research) and Dr Qing-Hai Fan (Biosecurity New Zealand) for reviews and comments. The author was supported by the New Zealand Foundation for Research, Science and Technology through backbone funding of the “Defining New Zealand’s Land Biota” programme.

References

- Adrain, J.M. (2011) Class Trilobita Walch, 1771. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 104–109.
- Ahyong, S.T., Lowry, J.K., Alonso, M., Bamber, R.N., Boxshall, G.A., Castro, P., Gerken, S., Karaman, G.S., Goy, J.W., Jones, D.S., Meland, K., Rogers, D.C. & Svavarsson, J. (2011) Order Scorpiones C.L. Koch, 1850. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 165–191.
- Arillo, A. & Engel, M.S. (2006) Rock crawlers in Baltic amber (Notoptera: Mantophasmatodea). *American Museum Novitates*, 3539, 1–10.
- Aristov, D.S. (2009) A new family of the order Grylloblattida (Insecta) from the Middle Permian of Russia. *Paleontologicheskii Zhurnal*, (2), 59–63.
- Beccaloni, G.W. & Eggleton, P. (2011) Order Blattodea Brunner von Wattenwyl, 1882. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 199–220.
- Bamber, R.N. (2011) Class Pycnogonida Latreille, 1810. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 110–111.
- Beaulieu, F., Dowling, A.P.G., Klompen, H., Moraes, G.J. de & Walter, D.E. (2011) Superorder Parasitiformes Reuter, 1909. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 123–128.
- Brock, P. & Marshall, J. (2011) Phasmida Leach, 1815. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 198–198.
- Chapman, A.D. (2009) Numbers of living species in Australia and the world. Second edition. – Australian Biodiversity Information Services, Toowoomba.
- Eberhard, M.J.B., Picker, M.D. & Klass, K.-D. (2011) Sympatry in Mantophasmatodea, with the description of a new species and phylogenetic considerations. *Organisms Diversity & Evolution*, 11(1), 43–59.
- Dunlop J.A. (2010) Geological history and phylogeny of Chelicerata. *Arthropod Structure & Development*, 39, 124–142.
- Dunlop J.A. & Penney, D. (2011) Order Araneae Clerck, 1757. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 149–153.
- Dunlop J.A., Penney D. & Jekel D. (2011) A summary list of fossil spiders and their relatives. *In*: Platnick, N. I. (ed.) The world spider catalog, version 12.0. American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html> DOI: 10.5531/db.iz.0001.
- Footitt, R & Adler, P. (eds) (2009) *Insect Biodiversity: Science and Society*. Blackwell Publishing, London, 623 pp.
- Giribet, G. & Edgecombe, G.D. (2012) Reevaluating the Arthropod Tree of Life. *Annual Review of Entomology*, 57, 167–186.
- Grimaldi, D. & Engel, M.S. (2005) *Evolution of the Insects*. Cambridge University Press, Cambridge, New York.
- Guglielmo, A.A., Robbins, R.G., Apaneskevich, D.A., Petney, T.N., Estrada-Peña, A., Horak, I.G., Shao, R. & Barker, S. (2010) The Argasidae, Ixodidae and Nuttalliellidae (Acari: Ixodida) of the world: a list of valid species names. *Zootaxa*, 2528, 1–28.
- Harvey, M.S. (2011) Order Pseudoscorpiones de Geer, 1778. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 119–120.
- Hong, Y.C. (2007) Discovery of the fossil glosselytrods (Insecta: Glosselytrodea) from Shaanxi, China. *Acta Entomologica Sinica*, 50(3), 271–280.
- Holzenthal, R.W., Morse, J.C. & Kjer, R.J., (2011) *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 209–211.
- Ingrisch, S. (2011) Order Orthoptera Oliver, 1789. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 195–197.
- Janssens, F. & Christiansen, K.A. (2011) Class Collembola Lubbock, 1870. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 192–194.
- Kury, A.B. (2011) Order Opiliones Sundevall, 1833. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 112–114.
- Mendes, L.F. (2002) Taxonomy of Zygentoma and Microryphia: historical overview, present status and goals for the new millennium. Proc. Xth International Colloquium on Apterygota, České Budějovice 2000: Apterygota at the Beginning of the Third Millennium. Elsevier GmbH.
- Minelli, A. (2011a) Class Trilobita, Class Symphyla and Class Pauropoda. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 157–158.
- Mound, L.A. (2011) Order Thysanoptera Haliday, 1836. *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148, 201–202.
- Nieukerken, E.J. van, Kaila, L., Kitching, I.J., Kristensen, N.P., Lees, D.C., Minet, J., Mitter, C., Mutanen, M., Regier, J.C., Simonsen, T.J., Wahlberg, N., Yen, S.-H., Zahiri, R., Adamski, D., Baixeras, J., Bartsch, D., Bengtsson, B.Å., Brown, J.W., Bucheli, S.R.,

- Davis, D.R., De Prins, J., De Prins, W., Epstein, M.E., Gentili-Poole, P., Gielis, C., Hättenschwiler, P., Hausmann, A., Holloway, J.D., Kallies, A., Karsholt, O., Kawahara, A.Y., Koster, S. (J.C.), Kozlov, M.V., Lafontaine, J.D., Lamas, G., Landry, J.-F., Lee, S., Nuss, M., Park, K.-T., Penz, C., Rota, J., Schintlmeister, A., Christian Schmidt, B., Sohn, J.-C., Solis, M.A., Tarmann, G.M., Warren, A.D., Weller, S., Yakovlev, R.V., Zolotuhin, V.V. & Zwick, A. (2011) Order Lepidoptera Linnaeus, 1758. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 212–221.
- Pape, T., Blagoderov, V. & Mostovski, M.B. (2011, this volume) Order Diptera Linnaeus, 1758. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 222–229.
- Prendini, L. (2011a) Order Scorpiones C.L. Koch, 1850. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 115–117.
- Prendini, L. (2011b) Solifugae Sundevall, 1833. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 118–118.
- Prendini, L. (2011c) Palpigradi Thorell, 1888. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 121–121.
- Prendini, L. (2011d) Ricinulei Thorell, 1876. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 122–122.
- Prendini, L. (2011e) Amblypygi Thorell, 1883. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 154–154.
- Prendini, L. (2011f) Thelyphonida Latreille, 1804. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 155–155.
- Prendini, L. (2011g) Schizomida Petrunkevitch, 1945. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 156–156.
- Prokop, J. & Nel, A. (2007) An enigmatic Palaeozoic stem-group: Paoliida, designation of new taxa from the Upper Carboniferous of the Czech Republic (Insecta: Paoliidae, Katerinkidae fam. n.). *African Invertebrates*, 48 (1), 77–86
- Rasnitsyn, A.P. & Quicke D.L.J. (eds) (2002) *History of Insects*. Kluwer Academic Publishers, Dordrecht, Boston, London, 517 pp.
- Selden, P.A., Shear, W.A. & Sutton, M.D. (2008) Fossil evidence for the origin of spider spinnerets, and a proposed arachnid order. *Proceedings of the National Academy of Sciences of the USA*, 105, 20781–20785.
- Schatz, H., Behan-Pelletier, V.M., OConnor, B.M. & Norton, R.A. (2011) Suborder Oribatida van der Hammen, 1968. *In: Zhang, Z.-Q. (Ed.), Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 141–148.
- Shcherbakov, D.E. (2011) The alleged Triassic palaeodictyopteran is a member of Titanoptera. *Zootaxa*, 3044, 65–68.
- Shear, W. (2011) Class Diplopoda de Blainville in Gervais, 1844. *In: Zhang, Z.-Q. (Ed.), Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 159–164.
- Shultz, J.W. (2007) A phylogenetic analysis of the arachnid orders based on morphological characters. *Zoological Journal of the Linnean Society*, 150, 221–265.
- Slipinski, S.A., Leschen, R.A.B. & Lawrence, J.F. (2011) Order Coleoptera Linnaeus, 1758. *In: Zhang, Z.-Q. (Ed.), Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 203–208.
- Szeptycki, A. (2007) Catalogue of the world Protura. Wydawnictwa Instytutu Systematyki i Ewolucji Zwierząt Polskiej Akademii Nauk, Kraków.
- Trautwein, M.D., Wiegmann, B.M., Beutel, R., Kjer, K.M. & Yeates, D.K. (2012) Advances in insect phylogeny at the dawn of the postgenomic era. *Annual Review of Entomology*, 57, 449–68.
- Trueman, J.W.H. (2007) A brief history of the classification and nomenclature of Odonata. *Zootaxa*, 1668, 381–394.
- Walter, D.E., Bolton, S., Uusitalo, M. & Zhang, Z.-Q. (2011) Suborder Endeostigmata Reuter, 1909. *In: Zhang, Z.-Q. (ed.) Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 139–140.
- Zhang, Z.-Q., Fan, Q.-H., Pesic, V., Smit, H., Bochkov, A.V., Khaustov, A.A., Baker, A., Wohltmann, A., Wen, T.-H., Amrine, J.W., Beron, P., Lin, J.-Z., Gabrys, G. & Husband, R. (2011) Order Trombidiformes Reuter, 1909. *In: Zhang, Z.-Q. (ed.) Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 129–138.

Author name and address

Zhi-Qiang Zhang, New Zealand Arthropod Collection, Landcare Research, Private Bag 92170, Auckland, New Zealand; zhangz@landcareresearch.co.nz