

NOTES ON SOME PARASITES OF SUGAR-CANE INSECTS
IN JAVA, WITH DESCRIPTIONS OF NEW *HYMEN-
OPTERA CHALCIDOIDEA*.*

By A. A. GIRAULT.

HERR P. VAN DER GOOT, Entomologist of the Experiment Station of the Java Sugar-cane Industry at Pasoeroean, Java, was kind enough to send to me for identification a number of egg-parasites of sugar-cane insects upon part of which I report on in the following pages :

1. *Gonatocerus bifasciativentris*, new species.

Female.—Length, 1.20 mm. Black and golden yellow, and belonging to the group of species with graceful fore wings; the abdomen as in *rivalis*, the ovipositor exerted for a length equal to a third that of the abdomen. Funicle and club, the propodeum, cephalic third or less of mesoscutum, cephalic half of parapside (making two triangular spots on each side), a subquadrate spot at base of scutellum at the meson, immediate base of abdomen, the exerted valves of the ovipositor and a broad black band just distad of centre of abdomen, in the dorsal aspect, sometimes narrowly divided into two stripes, velvety black. Pedicel yellow, suffused with dusky, the scape yellow, dusky along dorsal and ventral edges. Mesopleurum black. Fore wings with about twenty-one lines of discal cilia where broadest; marginal vein long for the genus; posterior wings with a paired line of discal cilia along each margin. Funicle of antennæ with no globular joints, all longer than wide; 1 and 2 subequal, smallest; 3 somewhat longer, subequal to the pedicel; 4, 5 and 6 subequal, longest; 7 only slightly shorter than 6, while 8 shortens.

Fore wings fumated slightly along distal margin.

From many specimens, $\frac{2}{3}$ = inch objective, 1 = inch optic, Bausch and Lomb.

Male.—The same, but the abdomen sometimes with three black stripes. Longest funicle joints nearly thrice their own width.

From eight specimens; the same magnification.

Though coloured somewhat like the Australian *cingulatus* and *comptei*, this species resembles in habitus *spinozai* and *bicolor* of Australia and *rivalis* of North America because of the more slender abdomen, the exerted ovipositor, and the absence of globate joints in the antennal funicle. But it is quite slender. Eight males and fifty-four females.

Habitat.—Java.

Host.—Eggs of a leaf-hopper embedded in the leaves of sugar-cane.

* Contribution No. 9, Entomological Laboratory, Bureau of Sugar Experiment Stations, Bundaberg, Queensland.

Types.—In the Queensland Museum, Brisbane, one male, sixteen females, on a slide in xylol balsam.

2. *Trichogramma minutum* (Riley).

Four males, eleven females of this species reared from the eggs of *Chilo infuscatellum*, Sn.

A New Genus of Omphaline Eulophidae.

Omphalini.

Parachrysocharis, new genus.

Female.—Like *Chrysocharis*, Foerster, but the post-marginal vein absent; antennæ slender, three funicle and club joints, third club joint terminating in a spur. Stigmal vein long. Thorax without grooves; the parapsidal furrows complete.

Male.—The same, but the funicle four-jointed, three club joints, ten antennal joints. Antennæ clothed with long, fine hairs.

Type.—*Parachrysocharis javensis*, new species.

3. *Parachrysocharis javensis*, new species.

Female.—Length, 1.25 mm. Black, the base of the abdomen rather broadly, the ventral half of the thorax and the legs golden yellow; venation and antennæ dusky yellow; face also mostly golden yellow. Funicle joints all longer than wide, subequal and each barely longer than the pedicel; scape and pedicel paler.

From nine specimens; the same magnification.

Male.—The same.

From three specimens, the same magnification.

Described from three males, nine females, reared from leaf-hopper eggs on the leaves of sugar-cane, the host probably *Flata affinis*; each egg holds one parasite. The hosts turn black.

Habitat.—Java.

Host.—*Flata affinis* (teste P. van der Goot).

Types.—In the Queensland Museum, Brisbane, one male, two females on a slide in xylol balsam.

4. *Cyrtogaster javensis*, new species.

Female.—Length, 1.90 mm. Dark metallic blue, the scutum reflecting greenish, wings hyaline; legs white and the antennæ yellowish, the latter with three ring-joints, each longer than the one preceding; the funicle joints subquadrate and more or less subequal. Post-marginal vein distinctly longer than the long stigmal. Coxæ concolorous. Parapsidal furrows half complete from cephalad. Punctate, the propodeum reticulated, with a median carina. Abdomen slender. Five funicle, three club joints, the club wider than the cylindrical funicle. Mandibles four-dentate, the outer tooth acute. Scutellum smoother toward apex.

From ten specimens; the same magnification.

Male.—Not known.

Described from ten females reared from the eggs of an unknown moth on the leaves of sugar-cane.

Habitat.—Java.

Host.—Lepidopterous eggs (probably Bombycidæ).

Types.—In the Queensland Museum, Brisbane, three females on tags, three pins and a slide with two others.

Runs to the genus in Ashmead's (1904) table.